Number of Carbohydrate Units

Monosaccharides = single unit
Disaccharides = two units
Oligiosaccharide = 3-10 units
Polysaccharide = 11+ units

Bonus:

- Can you name the most common Mono (4), Di(3), and Poly(4)-saccharides

Number of Carbons

3C = Triose
4C = Tetrose
5C = Pentose
6C = Hexose
7C = Heptose

Most common are 5 and 6 Carbon Carbohydrate

Bonus:

- Can you name the most common pentose?
- Can you name the most common hexoses?

Functional Group

Aldose = aldehyde



Ketose = ketone



Bonus:

- Can you name a common example of each?

D or L Isomer

- The orientation of the OH group furthest from the most oxidized end of a carbohydrate.
- The bottom OH on a properly drawn Fischer Projection



Size of Ring

Furanose = 5 member ring - Ald/Ket + OH 4 carbons a way



Pyranose = 6 member ring - Ald/Ket + OH 5 carbons away



Anomers

Definition:

- Diastereomers that differ in the configuration/orientation around the OH group on the carbon capable of mutarotation (hemiacetal or hemiketal carbon)
- BUDA (Beta Up, Down Alpha)



Epimers

Definition:
Two monosaccharide's that differ in the configuration around a single carbon.



D-allose



D-altrose



D-glucose



D-gulose



L-talose

Drawing Pyranose Rings

- Number the chain to decrease mistakes
- Left OH's Up
- Right OH's Down
- #6 - CH_2OH group up for D-isomers

Hemiacetal reaction





Drawing Furanose Rings





L-side OH are up

R-side OH are down

Drawing Disaccharides

- Formed by a dehydration reaction
- Draw a disaccharide given two monosaccharide's and the linkage
- Name disaccharides
- First ring (yl ending), Second normal

"Sucrose"
α-D-glucopyranosyl-(1,2)-
β-D-fructofuranose





"Lactose"
β -D-galactopyranosyl-(1,4)-α-D-glucopyranose

Hemiacetals, Acetals, Hemiketals, and Ketals



Hemiacetals and Hemiketals

- Capable of mutarotation
- React easily
- Reducing sugars







Acetals and Ketals

- Not Capable of mutarotation
- Not Reactive (hydrolysis)
- Not Reducing sugars

Monosaccharides

	4 Most Common Monosaccharide's			
Structural Isomers	D-Glucose	aldohexose	pyranose	bloodsugar, cellular respiration
	D-Galactose	aldohexose	pyranose	milk, yogurt, cell membranes
	D-Fructose	ketohexose	furanose	honey, sweetest sugar
	D-Ribose	aldopentose	furanose	DNA









Disaccharides

3 Most Common Disaccharides			
Maltose	α-D-Glucose + α-D-Glucose	α-1,4	beer, starch breakdown product
Lactose	β-D-Galactose + α-D-Glucose	β-1,4	milk sugar
Sucrose	α-D-Glucose + β-D-Fructose	α-β-1,2	table sugar

Polysaccharides

	4 Most Common Polysaccharide's			
Starch	Amylose	α-1,4	helix	plant energy storage
	Amylopectin	α-1,4 (main) α-1,6 (side)	treelike	plant energy storage
	Glycogen	similar to amylopectin	treelike	animal energy storage
	Cellulose	β-1,4	linear/ sheets	plant structural storage

Starch

Amylose:

- 25-1300 α-D-Glucose units
- α-1,4-glycosidic bonds
- Forms coils/helical/telephone structure
- Energy storage for plants





Amylopectin:

- 25-1300 α-D-Glucose units
- α-1,4-glycosidic bonds, branched every 25 glucose with a α-1,6-glycosidic bond
- Forms tree like structure
- Energy storage for plants



Glycogen

Glycogen:

- 25-1300 α-D-Glucose units
- α-1,4-glycosidic bonds, branched every 12-18 glucose with a α-1,6-glycosidic bond
- Forms tree like structure
- Similar to amylopectin, but more branched
- Energy storage for animals



Cellulose

Cellulose:

- 25-1300 β-D-Glucose units
- β-1,4-glycosidic bonds
- Forms linear chains, strong H-bonds leads to the formation of sheets
- Resistant to hydrolysis, indigestible by humans
- Most abundant organic substance in nature
- Chief structural component of plants and wooc



or





Mutarotation

- Process by which anomers are interconverted
- Equilibrium between cyclic and chain form.
- Occurs because hemiacetal carbon can open/close



Oxidation Reactions



Reduction Reaction



-"itol" acid

Kiliani-Fischer Reaction



Redox Tests

Redox Tests for Carbohydrates:

- Benedicts/Fehling/ Barfoeds – Cu^{+2} → Cu_2O (s) "Blue → Brick Red ppt
  - Benedicts/Fehling: general tests
  - Barfoeds: mono/di
- Tollens – Reduce Ag^{+} → Ag (s) "Silver Mirror"
- Sugar is Oxidized, Metals are Reduced

Functional Groups:

- Free Aldehydes
- α-hydroxyketones
- Hemiacetal





α-hydroxy ketone



Hemiacetal

Dehydration/Hydrolysis

Carbohydrate molecules are joined by Dehydration Reactions ($-H_2O$)
Di/Oligio/Polysaccharides are broken apart by Hydrolysis Reactions ($+H_2O$)



Dehydration/Hydrolysis

Carbohydrate molecules are joined by Dehydration Reactions (-H_2O)
Di/Oligio/Polysaccharides are broken apart by Hydrolysis Reactions (+H_2O)



Miscellaneous Applications

Sweeteners

TABLE 27.1 | Relative Sweetness of Sugars and Sugar Substitutes (based on fructose = 100)

Sugars	Relative sweetness	Sugar substitutes (common brand names)	Relative sweetness
Fructose	100	Sucralose (Splenda)	3.5×10^4
Invert sugar	75	Saccharin (Sweet 'N Low)	1.7×10^4
Sucrose	58	Acesulfame potassium (Sunette, Sweet One)	1.2×10^4
Glucose	43	Aspartame (Equal, NutraSweet)	1.0×10^4
Maltose	19	Rebiana (Truvia, PureVia)	1.2×10^4
Galactose	19	Neotame	4.1×10^5
Lactose	9.2		

Antigens / Blood Types